arXiv:1506.03444v1 [math.NT] 10 Jun 2015

Primality tests for specific classes of $N = k \cdot 2^m \pm 1$

Predrag Terzić

Bulevar Pera Ćetkovića 139 , Podgorica , Montenegro
e-mail: `pedja.terzic@hotmail.com`

Abstract: Polynomial time primality tests for specific classes of numbers of the form $k \cdot 2^m \pm 1$ are introduced .

Keywords: Primality test , Polynomial time , Prime numbers .
AMS Classification: 11A51 .

1 Introduction

In 1856 Edouard Lucas developed primality test for Mersenne numbers . The test was improved by Lucas in 1878 and Derrick Henry Lehmer in the 1930s , see [1] .In 1960 Kusta Inkeri provided unconditional , deterministic , lucasian type primality test for Fermat numbers , see [2] .In 1969 Hans Riesel formulated primality test , see [3] for numbers of the form $k \cdot 2^n - 1$ with k odd and $k < 2^n$. In 2006 Zhi-Hong Sun provided general primality criterion for numbers of the form $k \cdot 2^m \pm 1$,see [4] . In this note I present lucasian type primality tests for specific classes of $k \cdot 2^m \pm 1$.

Throughout this paper we use the following notations: $\mathbb{Z}$-the set of integers , $\mathbb{N}$-the set of positive integers , $\left(\frac{a}{p}\right)$-the Jacobi symbol , (m, n)-the greatest common divisor of m and n , $S_n(x)$-the sequence defined by $S_0(x) = x$ and $S_{k+1}(x) = (S_k(x))^2 - 2(k \geq 0)$.

2 Basic Lemmas and Theorems

Definition 2.1. For $P, Q \in \mathbb{Z}$ the Lucas sequence $\{V_n(P, Q)\}$ is defined by
$V_0(P, Q) = 2, V_1(P, Q) = P, V_{n+1}(P, Q) = PV_n(P, Q) - QV_{n-1}(P, Q)(n \geq 1)$
Let $D = P^2 - 4Q$. It is known that

$$V_n(P, Q) = \left(\frac{P+\sqrt{D}}{2}\right)^n + \left(\frac{P-\sqrt{D}}{2}\right)^n$$

Lemma 2.1. *Let $P, Q \in \mathbb{Z}$ and $n \in \mathbb{N}$. Then*

$$V_n(P, Q) = \sum_{r=0}^{[n/2]} \frac{n}{n-r} \binom{n-r}{r} P^{n-2r} (-Q)^r$$

Theorem 2.1. *(Zhi-Hong Sun)*

For $m \in \{2,3,4,...\}$ *let* $p = k \cdot 2^m \pm 1$ *with* $0 < k < 2^m$ *and* k *odd . If* $b, c \in \mathbb{Z}$ *,* $(p,c) = 1$ *and* $\left(\frac{2c+b}{p}\right) = \left(\frac{2c-b}{p}\right) = -\left(\frac{c}{p}\right)$ *then* p *is prime if and only if* $p \mid S_{m-2}(x)$*, where* $x = c^{-k}V_k(b,c^2) = \sum\limits_{r=0}^{(k-1)/2} \frac{k}{k-r}\binom{k-r}{r}(-1)^r(b/c)^{k-2r}$

For proof see Theorem 3.1 in [4] .

Lemma 2.2. *Let* n *be odd positive number , then*

$$\left(\tfrac{-1}{n}\right) = \begin{cases} 1, & \textit{if } n \equiv 1 \pmod 4 \\ -1, & \textit{if } n \equiv 3 \pmod 4 \end{cases}$$

Lemma 2.3. *Let* n *be odd positive number , then*

$$\left(\tfrac{2}{n}\right) = \begin{cases} 1, & \textit{if } n \equiv 1,7 \pmod 8 \\ -1, & \textit{if } n \equiv 3,5 \pmod 8 \end{cases}$$

Lemma 2.4. *Let* n *be odd positive number , then*
case 1. ($n \equiv 1 \pmod 4$*)*

$$\left(\tfrac{3}{n}\right) = \begin{cases} 1 & \textit{if } n \equiv 1 \pmod 3 \\ 0 & \textit{if } n \equiv 0 \pmod 3 \\ -1 & \textit{if } n \equiv 2 \pmod 3 \end{cases}$$

case 2. ($n \equiv 3 \pmod 4$*)*

$$\left(\tfrac{3}{n}\right) = \begin{cases} 1 & \textit{if } n \equiv 2 \pmod 3 \\ 0 & \textit{if } n \equiv 0 \pmod 3 \\ -1 & \textit{if } n \equiv 1 \pmod 3 \end{cases}$$

Proof. Since $3 \equiv 3 \pmod 4$ if we apply the law of quadratic reciprocity we have two cases . If $n \equiv 1 \pmod 4$ then $\left(\frac{3}{n}\right) = \left(\frac{n}{3}\right)$ and the result follows . If $n \equiv 3 \pmod 4$ then $\left(\frac{3}{n}\right) = -\left(\frac{n}{3}\right)$ and the result follows .

Lemma 2.5. *Let* n *be odd positive number , then*

$$\left(\tfrac{5}{n}\right) = \begin{cases} 1 & \textit{if } n \equiv 1,4 \pmod 5 \\ 0 & \textit{if } n \equiv 0 \pmod 5 \\ -1 & \textit{if } n \equiv 2,3 \pmod 5 \end{cases}$$

Proof. Since $5 \equiv 1 \pmod 4$ if we apply the law of quadratic reciprocity we have $\left(\frac{5}{n}\right) = \left(\frac{n}{5}\right)$ and the result follows .

Lemma 2.6. *Let* n *be odd positive number , then*
case 1. ($n \equiv 1 \pmod 4$*)*

$$\left(\frac{-3}{n}\right) = \begin{cases} 1 & \textit{if}\, n \equiv 1, 11 \pmod{12} \\ 0 & \textit{if}\, n \equiv 3, 9 \pmod{12} \\ -1 & \textit{if}\, n \equiv 5, 7 \pmod{12} \end{cases}$$

case 2. ($n \equiv 3 \pmod 4$*)*

$$\left(\frac{-3}{n}\right) = \begin{cases} 1 & \textit{if}\, n \equiv 5, 7 \pmod{12} \\ 0 & \textit{if}\, n \equiv 3, 9 \pmod{12} \\ -1 & \textit{if}\, n \equiv 1, 11 \pmod{12} \end{cases}$$

Proof.$\left(\frac{-3}{n}\right) = \left(\frac{-1}{n}\right)\left(\frac{3}{n}\right)$. Applying the law of quadratic reciprocity we have : if $n \equiv 1 \pmod 4$ then $\left(\frac{3}{n}\right) = \left(\frac{n}{3}\right)$. If $n \equiv 3 \pmod 4$ then $\left(\frac{3}{n}\right) = -\left(\frac{n}{3}\right)$. Applying the Chinese remainder theorem in both cases several times we get the result .

Lemma 2.7. *Let* n *be odd positive number , then*
case 1. ($n \equiv 1 \pmod 4$*)*

$$\left(\frac{7}{n}\right) = \begin{cases} 1 & \textit{if}\, n \equiv 1, 2, 4 \pmod 7 \\ 0 & \textit{if}\, n \equiv 0 \pmod 7 \\ -1 & \textit{if}\, n \equiv 3, 5, 6 \pmod 7 \end{cases}$$

case 2. ($n \equiv 3 \pmod 4$*)*

$$\left(\frac{7}{n}\right) = \begin{cases} 1 & \textit{if}\, n \equiv 3, 5, 6 \pmod 7 \\ 0 & \textit{if}\, n \equiv 0 \pmod 7 \\ -1 & \textit{if}\, n \equiv 1, 2, 4 \pmod 7 \end{cases}$$

Proof. Since $7 \equiv 3 \pmod 4$ if we apply the law of quadratic reciprocity we have two cases . If $n \equiv 1 \pmod 4$ then $\left(\frac{7}{n}\right) = \left(\frac{n}{7}\right)$ and the result follows . If $n \equiv 3 \pmod 4$ then $\left(\frac{7}{n}\right) = -\left(\frac{n}{7}\right)$ and the result follows .

Lemma 2.8. *Let* n *be odd positive number , then*
case 1. ($n \equiv 1 \pmod 4$*)*

$$\left(\frac{-6}{n}\right) = \begin{cases} 1 & \textit{if}\, n \equiv 1, 5, 19, 23 \pmod{24} \\ 0 & \textit{if}\, n \equiv 3, 9, 15, 21 \pmod{24} \\ -1 & \textit{if}\, n \equiv 7, 11, 13, 17 \pmod{24} \end{cases}$$

case 2. ($n \equiv 3 \pmod 4$*)*

$$\left(\frac{-6}{n}\right) = \begin{cases} 1 & \textit{if}\, n \equiv 7, 11, 13, 17 \pmod{24} \\ 0 & \textit{if}\, n \equiv 3, 9, 15, 21 \pmod{24} \\ -1 & \textit{if}\, n \equiv 1, 5, 19, 23 \pmod{24} \end{cases}$$

Proof. $\left(\frac{-6}{n}\right) = \left(\frac{-1}{n}\right)\left(\frac{2}{n}\right)\left(\frac{3}{n}\right)$. Applying the law of quadratic reciprocity we have : if $n \equiv 1 \pmod 4$ then $\left(\frac{3}{n}\right) = \left(\frac{n}{3}\right)$. If $n \equiv 3 \pmod 4$ then $\left(\frac{3}{n}\right) = -\left(\frac{n}{3}\right)$. Applying the Chinese remainder theorem in both cases several times we get the result .

Lemma 2.9. *Let n be odd positive number , then*

$$\left(\frac{10}{n}\right) = \begin{cases} 1 & \text{if } n \equiv 1,3,9,13,27,31,37,39 \pmod{40} \\ 0 & \text{if } n \equiv 5,15,25,35 \pmod{40} \\ -1 & \text{if } n \equiv 7,11,17,19,21,23,29,33 \pmod{40} \end{cases}$$

Proof. $\left(\frac{10}{n}\right) = \left(\frac{2}{n}\right)\left(\frac{5}{n}\right)$. Applying the law of quadratic reciprocity we have : $\left(\frac{5}{n}\right) = \left(\frac{n}{5}\right)$. Applying the Chinese remainder theorem several times we get the result .

3 The Main Result

Theorem 3.1. *Let $N = k \cdot 2^m - 1$ such that $m > 2$, $3 \mid k$, $0 < k < 2^m$ and*

$$\begin{cases} k \equiv 1 \pmod{10} \; with \; m \equiv 2,3 \pmod 4 \\ k \equiv 3 \pmod{10} \; with \; m \equiv 0,3 \pmod 4 \\ k \equiv 7 \pmod{10} \; with \; m \equiv 1,2 \pmod 4 \\ k \equiv 9 \pmod{10} \; with \; m \equiv 0,1 \pmod 4 \end{cases}$$

Let $b = 3$ and $S_0(x) = V_k(b,1)$, thus
N is prime iff $N \mid S_{m-2}(x)$

Proof. Since $N \equiv 3 \pmod 4$ and $b = 3$ from Lemma 2.2 we know that $\left(\frac{2-b}{N}\right) = -1$. Similarly , since $N \equiv 2 \pmod 5$ or $N \equiv 3 \pmod 5$ and $b = 3$ from Lemma 2.5 we know that $\left(\frac{2+b}{N}\right) = -1$. From Lemma 2.1 we know that $V_k(b,1) = x$. Applying Theorem 2.1 in the case $c = 1$ we get the result.

Theorem 3.2. *Let $N = k \cdot 2^m - 1$ such that $m > 2$, $3 \mid k$, $0 < k < 2^m$ and*

$$\begin{cases} k \equiv 3 \pmod{42} \; with \; m \equiv 0,2 \pmod 3 \\ k \equiv 9 \pmod{42} \; with \; m \equiv 0 \pmod 3 \\ k \equiv 15 \pmod{42} \; with \; m \equiv 1 \pmod 3 \\ k \equiv 27 \pmod{42} \; with \; m \equiv 1,2 \pmod 3 \\ k \equiv 33 \pmod{42} \; with \; m \equiv 0,1 \pmod 3 \\ k \equiv 39 \pmod{42} \; with \; m \equiv 2 \pmod 3 \end{cases}$$

Let $b = 5$ and $S_0(x) = V_k(b,1)$, thus
N is prime iff $N \mid S_{m-2}(x)$

Proof. Since $N \equiv 3 \pmod 4$ and $N \equiv 11 \pmod{12}$ and $b = 5$ from Lemma 2.6 we know that $\left(\frac{2-b}{N}\right) = -1$. Similarly , since $N \equiv 3 \pmod 4$ and $N \equiv 1 \pmod 7$ or $N \equiv 2 \pmod 7$ or $N \equiv 4 \pmod 7$ and $b = 5$ from Lemma 2.7 we know that $\left(\frac{2+b}{N}\right) = -1$. From Lemma 2.1 we know that $V_k(b, 1) = x$. Applying Theorem 2.1 in the case $c = 1$ we get the result.

Theorem 3.3. *Let* $N = k \cdot 2^m + 1$ *such that* $m > 2$, $0 < k < 2^m$ *and*

$$\begin{cases} k \equiv 1 \pmod{42} \textit{ with } m \equiv 2,4 \pmod 6 \\ k \equiv 5 \pmod{42} \textit{ with } m \equiv 3 \pmod 6 \\ k \equiv 11 \pmod{42} \textit{ with } m \equiv 3,5 \pmod 6 \\ k \equiv 13 \pmod{42} \textit{ with } m \equiv 4 \pmod 6 \\ k \equiv 17 \pmod{42} \textit{ with } m \equiv 5 \pmod 6 \\ k \equiv 19 \pmod{42} \textit{ with } m \equiv 0 \pmod 6 \\ k \equiv 23 \pmod{42} \textit{ with } m \equiv 1,3 \pmod 6 \\ k \equiv 25 \pmod{42} \textit{ with } m \equiv 0,2 \pmod 6 \\ k \equiv 29 \pmod{42} \textit{ with } m \equiv 1,5 \pmod 6 \\ k \equiv 31 \pmod{42} \textit{ with } m \equiv 2 \pmod 6 \\ k \equiv 37 \pmod{42} \textit{ with } m \equiv 0,4 \pmod 6 \\ k \equiv 41 \pmod{42} \textit{ with } m \equiv 1 \pmod 6 \end{cases}$$

Let $b = 5$ *and* $S_0(x) = V_k(b, 1)$ *, thus*
N *is prime iff* $N \mid S_{m-2}(x)$

Proof. Since $N \equiv 1 \pmod 4$ and $N \equiv 5 \pmod{12}$ and $b = 5$ from Lemma 2.6 we know that $\left(\frac{2-b}{N}\right) = -1$. Similarly , since $N \equiv 1 \pmod 4$ and $N \equiv 3 \pmod 7$ or $N \equiv 5 \pmod 7$ or $N \equiv 6 \pmod 7$ and $b = 5$ from Lemma 2.7 we know that $\left(\frac{2+b}{N}\right) = -1$. From Lemma 2.1 we know that $V_k(b, 1) = x$. Applying Theorem 2.1 in the case $c = 1$ we get the result.

Theorem 3.4. *Let* $N = k \cdot 2^m + 1$ *such that* $m > 2$, $0 < k < 2^m$ *and*

$$\begin{cases} k \equiv 1 \pmod 6 \textit{ and } k \equiv 1,7 \pmod{10} \textit{ with } m \equiv 0 \pmod 4 \\ k \equiv 5 \pmod 6 \textit{ and } k \equiv 1,3 \pmod{10} \textit{ with } m \equiv 1 \pmod 4 \\ k \equiv 1 \pmod 6 \textit{ and } k \equiv 3,9 \pmod{10} \textit{ with } m \equiv 2 \pmod 4 \\ k \equiv 5 \pmod 6 \textit{ and } k \equiv 7,9 \pmod{10} \textit{ with } m \equiv 3 \pmod 4 \end{cases}$$

Let $b = 8$ *and* $S_0(x) = V_k(b, 1)$ *, thus*
N *is prime iff* $N \mid S_{m-2}(x)$

Proof. Since $N \equiv 1 \pmod 4$ and $N \equiv 17 \pmod{24}$ and $b = 8$ from Lemma 2.8 we know that $\left(\frac{2-b}{N}\right) = -1$. Similarly , since $N \equiv 17 \pmod{40}$ or $N \equiv 33 \pmod{40}$ and $b = 8$ from Lemma 2.9 we know that $\left(\frac{2+b}{N}\right) = -1$. From Lemma 2.1 we know that $V_k(b, 1) = x$. Applying Theorem 2.1 in the case $c = 1$ we get the result.

References

[1] Crandall, Richard; Pomerance, Carl (2001) , ”Section 4.2.1: The Lucas-Lehmer test”, *Prime Numbers: A Computational Perspective* (1st ed.), Berlin: Springer, p. 167-170 .

[2] Inkeri, K., ”Tests for primality”, *Ann. Acad. Sci. Fenn.*, A I 279, 119 (1960).

[3] Riesel, Hans (1969) , ”Lucasian Criteria for the Primality of $N = h \cdot 2^n - 1$” , *Mathematics of Computation* (AmericanMathematical Society), 23 (108): 869-875 .

[4] Zhi-Hong Sun , ”Primality tests for numbers of the form $k \cdot 2^m \pm 1$” , *The Fibonacci Quarterly* , 44(2006), no.2, 121-130.